

Patient agreement to investigation or treatment

Haemorrhoidectomy

Authors: Colorectal Surgery

Brief description:

- Your surgeon has recommended that you undergo an operation to remove your haemorrhoids (piles). Surgery is required for certain types of haemorrhoids, or where other forms of treatment have not been able to control the symptoms.
- If you have further questions please contact our Colorectal Specialist Sister Karen Goodwin on 01223 217923 or speak to the doctor or Ward staff when you come to the hospital.
- Here, we explain some of the aims, benefits, risks and alternatives to this procedure (operation/treatment). We want you to be informed about your choices to help you to be fully involved in making any decisions.
- Please ask about anything you do not fully understand or wish to have explained in more detail.
- If you would like this information in another format or language or would like help completing the form, please ask a member of our staff.

Please bring this form with you to hospital

- You will be asked to read this form carefully, and you and your doctor (or other appropriate healthcare professional) will sign it to document your consent.
- All our consent forms are available on the Addenbrooke's website: <http://www.addenbrookes.org.uk/consent>
- Remember, you can change your mind about having the procedure at any time.

For staff use:

Does the patient have any special requirements? (For example, requires an interpreter or other additional communication method)

.....

.....

About haemorrhoidectomy

Haemorrhoids (piles) are small, blood-filled cushions located just inside the back passage (internal haemorrhoids) but can sometimes form outside the anus (external haemorrhoids). Problems with haemorrhoids are very common, but the symptoms often disappear without any treatment. Haemorrhoids may cause bleeding from the back passage, anal discomfort or itchiness, discharge of mucus, or prolapse with a lump coming through the anus. Very occasionally, blood clots inside a haemorrhoid and causes a painful hard lump on the outside of the anus (thrombosed external haemorrhoid). Risk factors for developing haemorrhoids include straining at stool, constipation, low fibre diets, obesity and pregnancy. The main ways to avoid having problems with haemorrhoids are to eat a high fibre diet and to avoid straining at stool.

Before your procedure

- Most patients attend a pre-admission clinic, when you will meet a member of the surgical team, usually a House Officer (junior doctor) or specialist nurse.
- At this clinic, we shall ask you for details of your medical history and carry out any necessary clinical examinations and investigations. This is a good opportunity for you to ask us any questions about the procedure, but please feel free to discuss any concerns you might have at any time.
- You will be asked if you are taking any tablets or other types of medication - these might be ones prescribed by a doctor or bought over the counter in a pharmacy. It helps us if you bring with you details of anything you are taking (for example bring the packaging with you).
- This procedure usually involves the use of general or regional anaesthesia. Very occasionally local anaesthesia with sedation is used. See below for further details about the types of anaesthesia/sedation we shall use.
- Most people who have this type of procedure have it performed as a day case procedure. You may need to stay overnight if you have other medical problems. Sometimes we can predict whether you will need to stay for longer than usual - your doctor will discuss this with you before you decide to have the procedure.
- You may be admitted on the day of your surgery or the day before. You should start to take a mild laxative such as lactulose (20mls twice a day) from two days prior to the operation. Just before surgery the nurse will give you an enema to empty the bowel.

Prevention of Surgical Site Infection (SSI)

It may be necessary during the procedure to shave another area of your body e.g. your thigh to allow attachment of a pad for the diathermy machine (used to seal blood vessels), so that the pad sticks to your skin to achieve the best and safest performance.

Hair removal from the site of the operation up to sixty minutes before surgery reduces the risk of infection. This means that the hair removal procedure is usually carried out on the operating table. You must not shave the area yourself; this will be carried out in the operating theatre. Shaving at home, or the night before surgery, increases the risk of infection as no matter how careful you are the skin may become irritated and this could increase the risk of infection.

During the procedure

- At the start of your procedure, you will be given the necessary anaesthetic and/or sedation - see below for details of this.
- Haemorrhoids usually have an external component outside the anus as well as the main component inside the anal canal. Depending on the number of piles, these are removed in two to four separate areas. The tissues may then be closed with a stitch or the wounds may be left open. Most of the stitch lies inside the anal canal but some will be on the outside. The stitch dissolves over a period of two to four weeks. No special dressings are required.

After the procedure

- If you have had a general anaesthetic you will wake up in the recovery room after your operation. You might have an oxygen mask on your face to help you breathe. You might also wake up feeling sleepy.
- After this procedure, most people will have a small, plastic tube in one of the veins of their arm. This might be attached to a bag of fluid (called a drip), which feeds your body with fluid until you are well enough to eat and drink by yourself.
- While you are in the recovery room, a nurse will check your pulse and blood pressure regularly. When you are well enough to be moved, you will be taken to a ward.
- Sometimes, people feel sick after an operation, especially after a general anaesthetic, and might vomit. If you feel sick, please tell a nurse and you will be offered medicine to make you more comfortable.
- **Eating and drinking:** You may eat and drink normally, and we recommend a high fibre diet and fluid intake of at least six to ten glasses of water daily.
- **Getting around and about:** Within a few hours of your operation, you will be encouraged to get up and walk around with assistance.
- **When you can leave hospital:** Discharge from hospital will be the same day (for planned daycase surgery) or the following day. You should expect to have your bowels open within two to three days and this will be uncomfortable at first. Please use the laxative you will be given to help reduce any discomfort on having your bowels open.
- **Special measures you need to take after the procedure:** A small amount of bleeding is expected. Over the first few weeks you may notice some change in your ability to control wind; this will resolve. Provided you feel comfortable, there are no restrictions on activity and you may lift, drive and go back to work.

- **Check-ups and results:** Follow up is not usually required. Before you leave hospital, you will be given details if you need to return to see us, for example outpatient clinics.

Pain relief

In order to minimise the pain associated with your operation, a number of measures will be taken:

- At the time of surgery, local anaesthetic will be injected. This will provide pain relief for much of the day.
- After surgery you will be given painkillers to take by mouth
- You will have a laxative for two weeks after the operation to prevent constipation.
- You may have sitz baths (a 15 minute bath in water as warm as you can tolerate) several times daily or as often as you require them. These are very soothing and provide several hours of pain relief.

Intended benefits of the procedure

- Haemorrhoidectomy removes troublesome haemorrhoids – including the external component/skin tags. It therefore controls both bleeding and prolapse.

Who will perform my procedure?

- Your surgery will be performed by a surgeon with appropriate experience – usually a Consultant or senior Specialist Registrar. A more junior surgeon in training may carry out the procedure but only under supervision of a more experienced surgeon.

Alternative procedures that are available

- Haemorrhoidectomy is generally only recommended when outpatient treatment has failed or is not suitable. There are a variety of other methods for surgical treatment of haemorrhoids but we do not favour any of these at present.

Serious or frequently occurring risks

Haemorrhoidectomy is generally a very safe operation with few risks, but, as with any surgical procedure, complications do occasionally occur; about 2% patients need to have a second anaesthetic to attend to a complication. Immediately after the operation, a few patients find it difficult to pass urine and a catheter may be required to empty the bladder. Around 5% of patients experience more bleeding than usual and this may need re-admission to hospital for observation or, rarely, another operation. Infection is very rare. The sutures that are placed sometimes separate, leaving an open wound: do not be concerned if this happens as healing will still take place. In the long term, recurrence of symptoms is rare although a few patients develop skin tags. Rarely, scarring can cause narrowing of the anus.

If you suffer from urinary symptoms due to large prostate you might be at increased risk of urinary problems after surgery.

Information and support

In the period following your operation you should contact your GP or the ward if you notice any of the following problems:

- Increasing pain, redness, swelling or discharge
- Severe bleeding
- Constipation for more than three days despite using a laxative
- Difficulty in passing urine
- High temperature over 38° or chills
- Nausea or vomiting

Your anaesthesia

Several different kinds of anaesthesia can be used, and the method will be tailored to your particular needs and wishes.

General anaesthesia

During general anaesthesia you are put into a state of unconsciousness and you will be unaware of anything during the time of your operation. Your anaesthetist achieves this by giving you a combination of drugs.

Before your operation

Before your operation your anaesthetist will visit you in the ward, although occasionally this will happen in a pre-anaesthetic assessment clinic. If you are a day case patient it may not be until just before your operation. The anaesthetist who looks after you on the day of your operation is the one who is responsible for making the final decisions about your anaesthetic. He or she will need to understand about your general health, any medication that you are taking and any past health problems that you have had. Your anaesthetist will want to know whether or not you are a smoker, whether you have had any abnormal reactions to any of the drugs or if you have any allergies. They will also want to know about your teeth, whether you wear dentures, have caps or a plate. Your anaesthetist needs to know all these things so that he or she can assess how to look after you in this vital period. Your anaesthetist may examine your heart and lungs and may also prescribe medication that you will be given shortly before your operation, the pre-medication or 'pre-med'.

Pre-medication is the name given to medication (drugs) given to you some hours before your operation. These drugs may be given as tablets, injections or liquids (to children). They relax you and may send you to sleep. They are not always given.

Do not worry if you do not have a pre-med, your anaesthetist has to take many factors into account in making this decision and will take account of your views on the topic if possible. Do not be worried about your anaesthetic. When your anaesthetist visits you before your operation, this is the time to ask all the questions that you may have, so that you can forget your fears and worries.

Before your operation you will usually be changed into a gown and wheeled to the operating suite into an anaesthetic room. This is an ante-room outside the theatre. The anaesthetist, his or her assistant and nurses are likely to be present. An intravenous line (drip) may be inserted. Monitoring devices may be attached to you, such as a blood pressure cuff or a pulse oximeter. A pulse oximeter is usually a little red light in a small box, which is taped to your finger. It shows how much oxygen you have in your blood and is one of the vital monitors that an anaesthetist uses during your operation to ensure that you remain in the best of health. You may be given some oxygen to breathe. It is common practice nowadays to allow a parent into the anaesthetic room with children: as the child goes unconscious, the parent will usually be asked to leave.

During your operation

While you are unconscious and unaware your anaesthetist remains with you at all times. He or she monitors your condition and administers the right amount of anaesthetic drugs to maintain you in the correct level of unconsciousness for the period of the surgery. Your anaesthetist is constantly aware of your condition and trained to respond. Your anaesthetist will be monitoring such factors as heart rate, blood pressure, heart rhythm, body temperature and breathing. He or she will also constantly watch your need for fluid or blood replacement. If you have any other medical conditions, your anaesthetist will know of these from your pre-operative assessment and be able to treat them during surgery.

After your operation

After your operation your anaesthetist continues to monitor your condition carefully. You will probably be transferred to a recovery ward where specially trained nurses, under the direction of anaesthetists, will look after you. Your anaesthetist and the recovery nurses will ensure that all the anaesthetic effects are reversed and that you are closely monitored as you return to full consciousness. You may be given some oxygen to breathe in the recovery area, and may find that intravenous drips have been inserted whilst you are unconscious in theatre and that these will be replacing fluids that you might require. You will be given medication for any pain that you might feel, and systems, such as Patient Controlled Anaesthesia (PCA) may be set up to continue pain control on the ward.

You are likely to feel drowsy and sleepy at this stage. Some patients feel sick, others may have a sore throat related to the insertion of the breathing tube during surgery.

During this time it is important that you relax as much as you can, breathe deeply, do not be afraid to cough, and do not hesitate to ask the nursing staff for any pain relief, and about any queries you may have. You are likely to have hazy memories of this time and some patients experience vivid dreams. Once you are fully awake you will be returned to the ward, and if you are a day patient will be allowed to go to the waiting area to fully recover before you are accompanied home. Do not expect to feel completely normal immediately!

What are the risks of general anaesthesia?

In modern anaesthesia, serious problems are uncommon. Risks cannot be removed completely, but modern equipment, training and drugs have made it a much safer procedure in recent years. The risk to you as an individual will depend on; whether you have any other illness, personal factors (such as smoking or being overweight) or surgery which is complicated, long or done in an emergency. Please discuss any pre-existing medical condition with your anaesthetist.

- Very common and common side effects (1 in 10 or 1 in 100 people)
Feeling sick and vomiting after surgery, sore throat, dizziness, blurred vision, headache, itching, aches, pains and backache, pain during injection of drugs, bruising and soreness, confusion or memory loss.
- Uncommon side effects and complications (1 in 1000 people)
Chest infection, bladder problems, muscle pains, slow breathing (depressed respiration), damage to teeth, lips or tongue, an existing medical condition getting worse, awareness (becoming conscious during your operation).
- Rare or very rare complications (1 in 10,000 or 1 in 100,000)
Damage to the eyes, serious allergy to drugs, nerve damage, death, equipment failure.

Spinal or caudal anaesthetic

- For some operations on the rectum/anus, a spinal anaesthetic can be used instead of a general anaesthetic. A spinal anaesthetic may be safer for some patients and be a more suitable anaesthetic than a general anaesthetic
- **What is a spinal?** A local anaesthetic is injected through a very fine needle into the small of your back. This will numb the nerves around your bottom and the back of your legs. Normally you will also have some light sedation so that you are not so aware of the operation. The amount of sedation can be adjusted so that you are not anxious and have reduced awareness without being unconscious. You should not feel any pain during the operation but you may be aware of other sensations.
- **Advantages of spinal anaesthesia**
There may be:
 - Less effect on the heart and lungs
 - Less sickness and vomiting
 - Excellent pain relief immediately after surgery
 - Less risk of injury when you are put into the position for your surgery

- **After your spinal:** You will return to the ward and can normally drink fluids and eat a light diet within an hour of the operation. You will remain in bed until you have full muscle power back in your legs. Please ask for help when you first get out of bed.
- As sensation returns you may experience some tingling in the skin as the spinal wears off. If you become aware of some pain from the operation site then you should take some pain relief. You should tell the ward staff about any concerns or worries that you have.

Side effects and complications:

- As with all anaesthetic techniques there is a possibility of unwanted side effects or complications.

Uncommon side effects include:

- Headache – When the spinal wears off and you begin to move around there is a risk of developing a headache.
- Difficulty passing water (urinary retention) – You may find it difficult to empty your bladder normally as long as the spinal lasts. Your bladder will work normally when the spinal has worn off.
- Pain during injection – Occasionally you may feel pain or ‘pins and needles’ in your legs or bottom during the injection. You should tell your anaesthetist immediately as this may indicate irritation or injury to a nerve and the needle will have to be repositioned.

Rare complications:

- Nerve damage – This is a rare complication of spinal anaesthesia. There may be temporary loss of sensation, pins and needles and sometimes muscle weakness that may last for a few days or even weeks but almost all of these make a full recovery in time. Permanent nerve damage is even more rare and has about the same chance of occurring as major complications of general anaesthesia.

Local anaesthesia

A third alternative is controlled **sedation** and injection of **local anaesthetic**.

In local anaesthesia the local anaesthetic drug is injected into the skin and tissues at the site of the operation. The area of numbness will be restricted and some sensation of pressure may be present, but there should be no pain. Local anaesthesia is used for minor operations such as stitching a cut. Usually a local anaesthetic will be given by the doctor doing the operation.

Sedation

Sedation is the use of small amounts of anaesthetic or similar drugs to produce a ‘sleepy-like’ state. It makes you physically and mentally relaxed during an investigation or procedure which may be unpleasant or painful (such as an endoscopy) but where your co-operation is needed.

The sedation is usually injected through a small needle or tube in the veins of the hand or in the arm. Sedation reduces the sensation of the injection of the local anaesthetic which usually stings.

You may remember a little about what happened but often you will remember nothing. This is known as 'conscious sedation', and may be used by other professionals as well as anaesthetists.

If you are having a regional or local anaesthetic, you may want to ask for some sedation as well.



We are currently working towards a smoke-free site. Smoking is only permitted in the designated smoking areas.

For advice and support in quitting, contact your GP or the free NHS Stop Smoking helpline on 0800 169 0 169

Help with this leaflet:



If you would like this information in another language, large print or audio format, please ask the department to contact Patient Information: 01223 216032 or

patient.information@addenbrookes.nhs.uk



Document history

Authors	Department of Colorectal Surgery
Department	Cambridge University Hospitals NHS Foundation Trust, Hills Road, Cambridge, CB2 0QQ www.cuh.org.uk
Contact number	01223 245151
Publish/Review date	February 2011/February 2014 (no changes made)
File name	General_surgery_haemorrhoidectomy
Version number/Ref	5/CF135

Consent Form (Adults)

Patient agreement to
investigation or treatment

For staff use only: Surname: First names: Date of birth: Hospital no: Male/Female: (Use hospital identification label)

Responsible health professional/job title

.....

Special requirements
(For example, other language/other communication method)

Name of proposed procedure or course of treatment

Haemorrhoidectomy

Statement of health professional

(To be filled in by a health professional with an **appropriate knowledge of the proposed procedure**, as specified in the Hospital's consent policy)

I have explained the procedure to the patient. In particular, I have explained:

- The intended benefits of the procedure
To control both bleeding and prolapse

- Any serious or frequently occurring risks from the procedures including those specific to the patient

Immediately after the operation, a few patients find it difficult to pass urine; More bleeding than usual; Recurrence of symptoms is rare although a few patients develop skin tags; Sutures that are placed sometimes separate, leaving an open wound; Scarring can cause narrowing of the anus.

- Any extra procedures that might become necessary during the procedure
- Blood transfusion Other procedure (please specify)

I have discussed what the treatment / procedure is likely to involve, the benefits and risks of any available alternative treatments (including no treatment) and any particular concerns of this patient.

- The following information leaflet has been provided: Haemorrhoidectomy
..... Version/Date/Ref: CF135, Version 5, February 2011

This procedure will involve:

- General and/or regional anaesthesia Local anaesthesia Sedation

Health professional's signature: Date:

Name (PRINT): Job title:

Contact details (if patient wishes to discuss details later)

- I have offered the patient information about the procedure but s/he has declined information.

Statement of the interpreter (if appropriate)

I have interpreted the information to the best of my ability, and in a way in which I believe s/he can understand:

Interpreter's signature..... Date:

Name (PRINT):

Important notes: (tick if applicable)

- The patient has withdrawn consent (ask patient to sign/date here)
- See also advance directive/living will (eg Jehovah's Witness form)

For staff use only: Surname: First names: Date of birth: Hospital no: Male/Female: (Use hospital identification label)

Statement of patient

Please read this form carefully. If your treatment has been planned in advance, you should already have your own copy, which describes the benefits and risks of the proposed treatment. If not, you will be offered a copy now. Do ask if you have any further questions. The staff at Addenbrooke's are here to help you. **You have the right to change your mind at any time before the procedure is undertaken, including after you have signed this form.**

Training doctors and other health professionals is essential to the continuation of the Health Service and improving the quality of care. Your treatment may provide an important opportunity for such training, where necessary under the careful supervision of a senior doctor. You may, however, decline to be involved in the formal training of medical and other students without this adversely affecting your care and treatment.

Please read the following:

I understand that I will have the opportunity to discuss the details of anaesthesia with an anaesthetist before the procedure, unless the urgency of my situation prevents this. (This only applies to patients having general or regional anaesthesia.)

I understand that you cannot give me a guarantee that a particular person will perform the procedure. The person undertaking the procedure will, however, have appropriate experience.

I understand that any procedure in addition to those described on this form will only be carried out if it is necessary to save my life or to prevent serious harm to my health.

I have been told about additional procedures which may become necessary during my treatment. I have listed below any procedures that **I do not wish, without further discussion, to be carried out.**

.....
I understand that any tissue (including blood) removed as part of the procedure or treatment will be anonymised and may be used for teaching or quality control, and stored or disposed of in a manner regulated by appropriate, ethical, legal and professional standards.

I understand that all research will be approved by a research ethics committee and undertaken in accordance with appropriate ethical, legal and professional standards.

I understand that the research may be conducted within a hospital, university, not for profit organisation or a company laboratory.

Please tick boxes to indicate you either agree/disagree to the three points below. Yes No

I agree that tissue (including blood) not needed for my own diagnosis or treatment can be used for **research which may include genetic research.** **If you wish** to withdraw your consent for the use of your tissue (including blood) for research, please contact the Patient Advice and Liaison Service at Addenbrooke's Hospital.

I agree to the use of photography for the purpose of diagnosis and treatment.

I agree to anonymised photographs being used for medical teaching.

I confirm that the risks, benefits and alternatives of this procedure have been discussed with me and I have read and understood the above and agree to the procedure (or course of treatment) on this form.

Patient's signature: **Date:**

Name (PRINT):

If the patient is unable to sign but has indicated his/her consent, a witness should sign below. Young people may also like a parent to sign here (see guidance notes).

Witness' signature: **Date:**

Name (PRINT):

Confirmation of consent (to be completed by a health professional when the patient is admitted for the procedure, if the patient has signed the form in advance)

On behalf of the team treating the patient, I have confirmed with the patient that s/he has no further questions and wishes the procedure to go ahead.

Signature **Date:**

Name (PRINT): **Job Title:**

Copy accepted by patient: yes / no (please circle)